

# Effects of Bio-Normalizer (A Food Supplementation) on Free Radical Production by Human Blood Neutrophils, Erythrocytes, and Rat Peritoneal Macrophages

JAMES AKIRA OSATO, PHD, LUDMILA G. KORKINA, PHD, DSC, LIBRADO A. SANTIAGO, PHD,  
AND IGOR B. AFANAS'EV, PHD, DSC

*From the Osato Research Institute, Gifu, Japan, and the Russian Institute of Pediatric Hematology, Moscow, Russia*

Date accepted: 15 July 1995

## ABSTRACT

Bio-normalizer, a natural Japanese health food prepared by the fermentation of *Carica papaya*, exhibits therapeutic properties against various pathologies including tumors and immunodeficiency. To understand the mechanism of bio-normalizer's therapeutic effects, we studied its action on the production of active oxygen species in cell-free systems (the Fenton reaction, the xanthine-xanthine oxidase system, and the hydrogen peroxide-hypochloride or hydrogen peroxide-horseradish peroxidase systems) and by human blood neutrophils and erythrocytes and rat peritoneal macrophages. Bio-normalizer efficiently inhibited the formation of oxygen radicals in cell-free systems and partly decreased spontaneous and radical release by dormant and activated phagocytes (neutrophils and macrophages). We suggest that bio-normalizer is able to enhance the intracellular production of innocuous superoxide ion and, at the same time, to diminish the formation of reactive hydroxyl radicals, perhaps by the inactivation of ferrous ions, the catalysts of the superoxide-driven Fenton reaction. We also propose that the normalization of an organism's superoxide level is one of the molecular mechanisms of bio-normalizer activity. *Nutrition* 1995;11:568-572

Key words: bio-normalizer, free radicals, neutrophils, macrophages

## INTRODUCTION

Bio-normalizer, a traditional natural Japanese health food prepared by the fermentation of *Carica papaya*, exhibits therapeutic effects in various human diseases. It was previously proposed that bio-normalizer efficiency in the treatment and prevention of severe pathologies is linked to its free radical scavenging properties such as the scavenging of hydroxyl radicals<sup>1</sup> and the inhibition of lipid peroxidation.<sup>2</sup> There is now a large body of evidence that free radical scavengers and antioxidants may inhibit neoplastic events at cellular and molecular levels,<sup>3</sup> prevent various environment-associated pathologies,<sup>4</sup> and diminish other disorders mediated by free radicals. It is also well established that the most important source of reactive oxygen species in mammals is phagocytosing cells such as circulating neutrophils and tissue macrophages.

Therefore, we have studied for the first time the effects of bio-normalizer on oxygen radical production by human blood neutrophils and erythrocytes and by rat peritoneal macrophages. We found that bio-normalizer both inhibited and stimulated the release of oxygen radicals by phagocytes, which is supposed to be an origin of bio-normalizer's in vivo biological activity. In addition, we showed that bio-normalizer scavenged free radicals produced in some cell-free systems.

## MATERIALS AND METHODS

### Chemicals

Bio-normalizer was produced by Sun-O International Inc., Gifu, Japan, by the fermentation of *Carica papaya* tropical herbal plants. A major component of bio-normalizer is a mixture of plant and yeast glycopolysaccharides. In addition, bio-

Correspondence to: Igor B. Afanas'ev, PhD, Vitamin Research Institute, Nauchny pr. 14A, Moscow 117820, Russia.




FIG. 1. Effect of soluble bio-normalizer (SBN) on the production of active oxygen species in cell-free systems:  $\diamond$ , the Fenton reaction ( $\text{H}_2\text{O}_2 + \text{FeSO}_4$ ), every point is an average of four experiments;  $\Delta$ , hydrogen peroxide-hypochloride system, every point is an average of three experiments;  $\circ$ , hydrogen peroxide-horseradish peroxidase system, every point is an average of five experiments; and  $\blacksquare$ , xanthine-xanthine oxidase system, every point is an average of four experiments. Experimental details are presented in the MATERIALS AND METHODS section. CL, chemiluminescence.

normalizer contains amino acids (for example, tryptophan, leucine, and glutamic and aspartic acids), vitamins B-6 and C, glucose, and the protease-antiprotease system of papain. Under our experimental conditions, the suspension of bio-normalizer granules was prepared daily in Hanks' balanced salt solution (HBSS). In most experiments, bio-normalizer particles were sedimented by centrifugation after a 2-h incubation in HBSS, and clear supernatant containing the soluble bio-normalizer (SBN) fraction was used in experiments. SBN contained all the above cited compounds excluding insoluble, high molecular weight glycopolysaccharides.

Cytochrome *c*, NADPH, lucigenin, luminol, menadione, phorbol myristate acetate (PMA), zymosan A, latex particles (1  $\mu\text{m}$ ), Ficoll-Hypaque, Monoprep, and bovine superoxide dismutase (SOD, EC 1.15.1.1) were from Sigma Chemical Co. (St. Louis, USA). Dextran sulfate and sodium metrizoate were from Pharmacia (Uppsala, Sweden).

#### Blood Drawing and Cell Isolation

Venous blood (2 ml) was obtained from healthy subjects after their informed consent. Heparin-treated blood was carefully layered on a dextran-metrizoate mixture (2 ml, 2:5 vol:vol) and incubated at room temperature for 30 min. Then, the leukocyte-rich top layer was placed on 1 ml Ficoll-Hypaque ( $d = 1.119$ ) and centrifuged at  $400 \times g$  for 30 min. The neutrophil-containing pellet was washed twice in cold HBSS and resuspended in a minimal essential medium (MEM). Cell suspensions contained  $>95\%$  neutrophils as measured by differential cell counts with a

Coulter counter. Neutrophil viability was 90–98% as assessed by the exclusion of 0.2% trypan blue dye.

Erythrocytes were collected from the bottom layer after dextran-metrizoate sedimentation, washed 3 times with 0.1 mol potassium-sodium phosphate buffer/L (pH 7.3), diluted with this buffer to the 2% hematocrit, and stored at  $4^\circ\text{C}$  until analyzed. Cells were counted by the Coulter method.

#### Animal and Macrophage Preparation

Adult male Wistar rats were injected intraperitoneally with 10 ml HBSS and killed 5 min later. The peritoneal lavage fluid was removed, placed on top of Monoprep solution ( $d = 1.068$ ), and centrifuged at  $400 \times g$  for 30 min. The upper portion containing mononuclear cells was washed twice with cold calcium- and magnesium-free HBSS and resuspended in MEM supplemented with 1% heat-inactivated calf serum. A differential cell count was performed microscopically using Giemsa staining: about 80% of cells were macrophages and monocytes and the others were lymphocytes, mast cells, and erythrocytes;  $>95\%$  of cells excluded trypan blue.

#### Measurement of the Production of Active Oxygen Species

The production of active oxygen species in the Fenton reaction and by the xanthine-xanthine oxidase, hydrogen peroxide-horseradish peroxidase (HRP), and hydrogen peroxide-hypochloride systems was measured by luminol- and lucigenin-amplified chemiluminescence in the chemiluminescence unit of an LKB luminometer (Model 1251, Wallan, Finland) at  $25^\circ\text{C}$ .

For the Fenton reaction,  $\text{FeSO}_4$  (2 mmol/L) and luminol (0.1 mmol/L) were incubated with or without bio-normalizer in 0.125 mol phosphate buffer/L (pH 7.4). Next, the reaction was started by adding hydrogen peroxide (1 mmol/L) and the chemiluminescence intensity was registered continuously.

For measurement in the xanthine-xanthine oxidase system, xanthine (0.1 mol/L) and lucigenin (10  $\mu\text{mol/L}$ ) were incubated with or without bio-normalizer in 0.125 mol  $\text{Na}_2\text{CO}_3$ /L (pH 7.8) and the reaction was started by adding xanthine oxidase (0.5 U/mg). In the hydrogen peroxide-HRP and hydrogen peroxide-hypochloride systems, luminol (0.1 mmol/L) and HRP (0.12 mg) or  $\text{NaClO}$  (1 mmol/L) were incubated with or without bio-normalizer in 0.01 mol Tris-HCl buffer/L and the reaction was started by adding hydrogen peroxide.

For the measurement of superoxide production by erythrocytes, an erythrocyte suspension (20  $\mu\text{l}$ ) and cytochrome *c* (50  $\mu\text{mol/L}$ ) were incubated with or without bio-normalizer in 0.1 mol potassium phosphate buffer/L (2 ml) with or without menadione (10  $\mu\text{mol/L}$ ) at room temperature for 60 min. After the sample was centrifuged, the light absorbance of the clear supernatant was measured at 550 nm on a Beckman DU 600 spectrophotometer (St. Louis, MO, USA). Superoxide content was calculated with the extinction coefficient of reduced cytochrome *c* equal to  $21000 \text{ mol} \cdot \text{L}^{-1} \cdot \text{cm}^{-1}$ .

To measure superoxide production by neutrophils and macrophages and NADPH oxidase activity,<sup>5</sup> we incubated a cell suspension ( $5 \times 10^5$  cells) and cytochrome *c* (50  $\mu\text{mol/L}$ ) with or without bio-normalizer in HBSS and continuously recorded the light absorbance at 550 nm. The reaction was started by adding PMA (10 ng/mL), stopped at the maximal rate by the addition of 0.05% Triton X-100, and restored 1 min later by the addition of NADPH (250  $\mu\text{mol/L}$ ).

The production of active oxygen species by neutrophils and macrophages was measured by the chemiluminescence method with lucigenin as a specific chemiluminescence probe for the detection of superoxide ion<sup>6</sup> and luminol as the




FIG. 2. Effects of bio-normalizer and soluble bio-normalizer (SBN) on luminol-amplified chemiluminescence (CL) of rat peritoneal macrophages. Every point is an average of four experiments. ●, bio-normalizer; □ or ■, SBN. A: Spontaneous CL. Basal CL is 20–70 mV/10<sup>6</sup> cells. The stimulator's effect of 1.25 mg bio-normalizer/ml is significantly different from control ( $p < 0.05$ ). B: CL activated by phorbol myristate acetate. The basal CL level is 250–400 mV/10<sup>6</sup> cells. C: Zymosan-activated CL. Basal CL level is 700–800 mV/10<sup>6</sup> cells. D: Latex-activated CL. Basal CL is 1100–1400 mV/10<sup>6</sup> cells. In all cases, the inhibitory effects of 5–20 mg bio-normalizer/ml or SBN are significantly different from control ( $p < 0.01$ ).

chemiluminescence probe for detecting hydroxyl and hydroxyl-like free radicals.<sup>7</sup> Cells ( $1 \times 10^5$ ) with or without bio-normalizer were mixed with HBSS (0.9 ml) containing 50  $\mu$ mol/L luminol or lucigenin. The cell-containing mixture was incubated at 37°C and mixed continuously for 5 min and spontaneous chemiluminescence was registered. Then, 50  $\mu$ l of activator (PMA, opsonized zymosan, or latex particles) was added and the maximal chemiluminescence response to the activator was measured.

#### Statistics

All measurements were performed in triplicate, and the results are expressed as an average  $\pm$  SD from three to four independent experiments.

#### RESULTS

SBN (Fig. 1) and bio-normalizer (data not shown) inhibited the production of oxygen radicals in a concentration-dependent manner in the cell-free systems studied. Thus, SBN  $I_{50}$  values were equal to 0.4, 4.5, 2.0, and 2.5 mg/ml for the Fenton reaction, and the hydrogen peroxide–hypochloride, hydrogen peroxide–HRP, and xanthine–xanthine oxidase systems, respectively. At the same time, SBN and bio-normalizer showed

only weak inhibitory activities in erythrocyte suspension with or without menadione (data not shown).

Bio-normalizer (but not SBN) slightly stimulated spontaneous luminol-amplified chemiluminescence of rat peritoneal macrophages at 0.2–1.25 mg/ml (Fig. 2A). The stimulatory activity of bio-normalizer particles increased after opsonization with autologous serum. Similar effects were observed with nonopsonized and opsonized zymosan. On the other hand, bio-normalizer and SBN inhibited luminol-amplified chemiluminescence of macrophages after their activation with soluble (PMA) or particular (latex or zymosan) stimuli (Fig. 2, B, C, and D). When lucigenin was applied as a chemiluminescence amplifier, bio-normalizer and SBN stimulated spontaneous, PMA, and zymosan-activated chemiluminescence (Fig. 3, A, B, and C) and inhibited latex-activated chemiluminescence at concentrations  $>1$  mg/ml (Fig. 3D). In all cases, lucigenin-amplified chemiluminescence was completely inhibited by SOD (10  $\mu$ g/ml). Similar results were obtained in the experiments with human neutrophils (data not shown).

SBN and bio-normalizer inhibited SOD-sensitive cytochrome *c* reduction by PMA-stimulated neutrophils ( $I_{50} = 10$  mg/ml) and slightly (not more than by 25%) decreased neutrophilic NADPH oxidase activity (data not shown).




FIG. 3. Effects of bio-normalizer and soluble bio-normalizer (SBN) on lucigenin-amplified chemiluminescence (CL) of rat peritoneal macrophages. Every point is an average of four experiments. ●, bio-normalizer; □ or ■, SBN. A: Spontaneous CL. Basal CL level is 2–10 mV/10<sup>6</sup> cells. The stimulatory effects of 5 mg bio-normalizer/ml and 20 mg SBN/ml are significantly different from control ( $p < 0.01$  and 0.05, respectively). B: CL activated by phorbol myristate acetate. Basal CL level is 25–60 mV/10<sup>6</sup> cells. The stimulatory effects of 5 mg bio-normalizer/ml or SBN are significantly different from control ( $p < 0.05$ ). C: Zymosan-activated CL. Basal CL level is 30–80 mV/10<sup>6</sup> cells. The stimulatory effect of 20 mg bio-normalizer/ml is significantly different from control ( $p < 0.05$ ). D: Latex-activated CL. Basal CL level is 100–150 mV/10<sup>6</sup> cells. The inhibitory effects of 5 mg bio-normalizer/ml bio-normalizer or SBN are significantly different from control ( $p < 0.05$ ).

#### DISCUSSION

Bio-normalizer is a complex mixture of various biologically active compounds. Therefore, it is not surprising that bio-normalizer can affect oxygen radical production by different pathways. We found that bio-normalizer inhibited the formation of hydroxyl radicals by the Fenton reaction, superoxide production by xanthine oxidase, and the formation of other active oxygen species by the hydrogen peroxide–hypochloride and hydrogen peroxide–HRP systems (Fig. 1). However, in contrast with cell-free systems, at the cellular level, bio-normalizer exhibited both stimulatory and inhibitory activities. Although the inhibitory effects of bio-normalizer may be explained by its scavenging properties, the nature of the stimulatory effect of bio-normalizer depends on different mechanisms.

The stimulation of spontaneous chemiluminescence by 0.2–1.25% bio-normalizer (Fig. 2A) is apparently a consequence of the particular receptor activation of phagocytes. A more complicated mechanism is possibly responsible for the stimulatory effects of bio-normalizer on lucigenin-amplified chemiluminescence (Fig. 3). We propose that this effect may originate from the interaction of macrophages with T lymphocytes. Macrophages primed with yeast polysaccharides can interact with T lymphocytes, thus enhancing interferon- $\gamma$  production.<sup>8</sup> Interferon- $\gamma$ , in turn, is able to prime stimulated

macrophages, thus increasing oxygen radical release.<sup>9</sup> Bio-normalizer contains a relatively large amount of yeast particles and yeast cell wall components; these components may be responsible for the stimulatory effect of bio-normalizer on superoxide production (measured via lucigenin-amplified chemiluminescence) by macrophages (Fig. 3, excluding latex-activated chemiluminescence; the reason of this exclusion remains unknown). On the other hand, the scavenging of hydroxyl radicals (measured via luminol-amplified chemiluminescence) by antioxidative bio-normalizer components is probably more important than the stimulation of their formation. This theory explains the fact that bio-normalizer inhibited luminol-amplified chemiluminescence (Fig. 2).

The ability of bio-normalizer to inhibit hydroxyl radical production and enhance superoxide production by phagocytes suggests the conversion of highly reactive hydroxyl radicals to innocuous superoxide ions. This process probably takes place in the cytoplasm or cellular membranes and possibly plays an important role in prevention of the destruction of biological molecules and structures.<sup>10</sup> It can also be a molecular basis of bio-normalizer normalizing activity because the optimal level of superoxide ion in the blood and tissues is thought to be an important factor of normal functioning of the immune system<sup>11</sup> and gene expression.<sup>12</sup>

## REFERENCES

1. Santiago LA, Osato JA, Hiramatsu M, et al. Free radical scavenging action of Bio-catalizer  $\alpha$ -p no.11 (Bio-normalizer) and its by-product. *Free Radic Biol Med* 1991;11:379
2. Santiago LA, Osato JA, Liu J, et al. Age-related increase in superoxide dismutase and thiobarbituric acid-reactive substances: effect of Bio-normalizer in aged rat brain. *Neurochem Res* 1993;18:711
3. Perchellet JP, Perchellet EM. Antioxidants and multistage carcinogenesis in mouse skin. *Free Radic Biol Med* 1989;7:377
4. Ames BN. Dietary carcinogens and anticarcinogens, oxygen radicals and degenerative disease. *Science* 1983;221:1256
5. Bellavite P, Jones OTC, Cross AR, et al. Composition of partially purified NADPH oxidase from pig neutrophils. *Biochem J* 1984;223:639
6. Gyllenhammar H. Lucigenin chemiluminescence in the assessment of neutrophil superoxide production. *J Immunol Methods* 1987;97:209
7. Allen RC, Loose LD. Phagocytic activation of a luminol dependent chemiluminescence in rabbit peritoneal macrophages. *Biochem Biophys Res Commun* 1976;69:245
8. Adams DO, Hamilton TA. Phagocytic cells: cytotoxic activities of macrophages. In: Galin JI, Goldstein IM, Snyderman RI, eds. *Inflammation: basic principles and clinical correlates*. New York:Raven Press, 1988:471
9. Ito M, Karmali R, Krim M. Effect of interferon on chemiluminescence and hydroxyl radical production in murine macrophages stimulated by PMA. *Immunology* 1985;56:533
10. Korkina LG, Afanas'ev IB, Samochatova EV, et al. Release of active oxygen radicals by leukocytes of Fanconi anemia patients. *J Leukoc Biol* 1992;52:357
11. Wing EJ, Ampel NM, Waheed A, et al. Macrophage colony-stimulating factor (M-CSF) enhances the capacity of murine macrophages to secrete oxygen reduction products. *J Immunol* 1985;135:2052
12. Oberley CW, Buetner GR. Role of superoxide dismutase in cancer: a review. *Cancer Res* 1979;39:1140

